

PRODUCTS OF k ATOMS IN KRULL MONOIDS

YUSHUANG FAN AND QINGHAI ZHONG

ABSTRACT. Let H be a Krull monoid with finite class group G such that every class contains a prime divisor. For $k \in \mathbb{N}$, let $\mathcal{U}_k(H)$ denote the set of all $m \in \mathbb{N}$ with the following property: There exist atoms $u_1, \dots, u_k, v_1, \dots, v_m \in H$ such that $u_1 \cdots u_k = v_1 \cdots v_m$. It is well-known that the sets $\mathcal{U}_k(H)$ are finite intervals whose maxima $\rho_k(H) = \max \mathcal{U}_k(H)$ depend only on G . If $|G| \leq 2$, then $\rho_k(H) = k$ for every $k \in \mathbb{N}$. Suppose that $|G| \geq 3$. An elementary counting argument shows that $\rho_{2k}(H) = kD(G)$ and $kD(G) + 1 \leq \rho_{2k+1}(H) \leq kD(G) + \lfloor \frac{D(G)}{2} \rfloor$ where $D(G)$ is the Davenport constant. In [11] it was proved that for cyclic groups we have $kD(G) + 1 = \rho_{2k+1}(H)$ for every $k \in \mathbb{N}$. In the present paper we show that (under a mild condition on the Davenport constant) for every noncyclic group there exists a $k^* \in \mathbb{N}$ such that $\rho_{2k+1}(H) = kD(G) + \lfloor \frac{D(G)}{2} \rfloor$ for every $k \geq k^*$. This confirms a conjecture of A. Geroldinger, D. Gryniewicz, and P. Yuan in [13].

1. INTRODUCTION

Let H be an atomic monoid. If an element $a \in H$ has a factorization $a = u_1 \cdots u_k$ into atoms $u_1, \dots, u_k \in H$, then k is called the length of the factorization, and the set $\mathsf{L}(a)$ of all possible lengths is called the set of lengths of a . For $k \in \mathbb{N}$, let $\mathcal{U}_k(H)$ denote the set of all $m \in \mathbb{N}$ with the following property: There exist atoms $u_1, \dots, u_k, v_1, \dots, v_m \in H$ such that $u_1 \cdots u_k = v_1 \cdots v_m$. Thus $\mathcal{U}_k(H)$ is the union of all sets of lengths containing k . The sets $\mathcal{U}_k(H)$ are one of the most investigated invariants in factorization theory which were introduced by S.T. Chapman and W.W. Smith in Dedekind domains ([7]). Their suprema $\rho_k(H) = \sup \mathcal{U}_k(H)$ were first studied in the 1980s for rings of integers in algebraic number fields ([8, 19]). Since then these invariants have been studied in a variety of settings, including numerical monoids, monoids of modules, noetherian and Krull domains (for a sample out of many we refer to [10, 4, 3, 15, 1]).

In the present paper we focus on Krull monoids with class group G such that every class contains a prime divisor. If $|G| \leq 2$, then $\mathcal{U}_k(H) = \{k\}$ and if G is infinite, then $\mathcal{U}_k(H) = \mathbb{N}_{\geq 2}$ for all $k \in \mathbb{N}$. Suppose that G is finite with $|G| \geq 3$. This setting includes holomorphy rings in global fields. For more examples we refer to [13], and a detailed exposition of Krull monoids can be found in [18, 14].

The unions $\mathcal{U}_k(H) \subset \mathbb{N}$ are finite intervals, say $\mathcal{U}_k(H) = [\lambda_k(H), \rho_k(H)]$, whose minima $\lambda_k(H)$ can be expressed in terms of $\rho_k(H)$ ([12, Chapter 3]). Elementary counting arguments (e.g. [14, Section 6.3]) show that, for every $k \in \mathbb{N}$, we have $\rho_{2k}(H) = kD(G)$ and that

$$(1.1) \quad kD(G) + 1 \leq \rho_{2k+1}(H) \leq kD(G) + \left\lfloor \frac{D(G)}{2} \right\rfloor.$$

Based on the Savchev-Chen Structure Theorem [17, Section 11.3] (resp. on a related result on the index of sequences) Gao and Geroldinger [11] showed that for every cyclic group G and every $k \in \mathbb{N}$ we have

2010 *Mathematics Subject Classification.* 11B30, 11R27, 13A05, 13F05, 20M13.

Key words and phrases. non-unique factorizations, sets of lengths, Krull monoids, zero-sum sequences.

Main part of this manuscript was written while the first author visited University of Graz. She would like to gratefully acknowledge the kind hospitality from the host institute and the authors would like to thank Professor Alfred Geroldinger for his many helpful suggestions. This work was supported by the Austrian Science Fund FWF (Project No. M1641-N26), NSFC (grant no. 11401542), and the Fundamental Research Funds for the Central Universities (grant no. 2652014033).

$\rho_{2k+1}(H) = kD(G) + 1$. In [13, Conjecture 3.3], the authors conjectured that for every noncyclic group G there exists a $k^* \in \mathbb{N}$ such that

$$\rho_{2k+1}(H) = kD(G) + \left\lfloor \frac{D(G)}{2} \right\rfloor \quad \text{for every } k \geq k^*.$$

We confirm this conjecture for wide classes of groups. For a precise formulation of our main result we need one more definition. Suppose that $G \cong C_{n_1} \oplus \dots \oplus C_{n_r}$ where $r, n_1, \dots, n_r \in \mathbb{N}$ with $1 < n_1 \mid \dots \mid n_r$, and set

$$D^*(G) = 1 + \sum_{i=1}^r (n_i - 1).$$

It is well-known that $D^*(G) \leq D(G)$. Equality holds for p -groups, groups of rank at most two, and others (see [12, Corollary 4.2.13], [5] for recent progress), but it does not hold in general ([16]). Here is our main result.

Theorem 1.1. *Let H be a Krull monoid with finite noncyclic class group G such that every class contains a prime divisor. Then there exists a $k^* \in \mathbb{N}$ such that*

$$\rho_{2k+1}(H) \geq (k - k^*)D(G) + k^*D^*(G) + \left\lfloor \frac{D^*(G)}{2} \right\rfloor \quad \text{for every } k \geq k^*.$$

In particular, if $D(G) = D^*(G)$, then

$$\rho_{2k+1}(H) = kD(G) + \left\lfloor \frac{D(G)}{2} \right\rfloor \quad \text{for every } k \geq k^*.$$

In [13], Geroldinger, Gryniewicz, and Yuan gave a list of groups for which the above result holds with $k^* = 1$. Furthermore, they showed that if $G \cong C_m \oplus C_{mn}$ with $n \geq 1$ and $m \geq 2$, then the result holds with $k^* = 1$ if and only if $n = 1$ or $m = n = 2$. It remains a challenging task to determine, for a given group G , the smallest possible $k^* \in \mathbb{N}$ for which the above statement holds.

It is well-known that the invariants $\rho_k(H)$ can be studied in an associated monoid of zero-sum sequences and this allows to use methods from Additive Combinatorics (see Lemma 2.1). In Section 2 we fix our notation and terminology. At the beginning of Section 3 we introduce our main concept in Definition 3.1 and after that we discuss the strategy of the proof.

2. PRELIMINARIES

Let \mathbb{N} denote the set of positive integers and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For real numbers $a, b \in \mathbb{R}$, we denote by $[a, b] = \{x \in \mathbb{Z} \mid a \leq x \leq b\}$ the discrete interval. For $n \in \mathbb{N}$ we denote by C_n a cyclic group of order n . Let G be a finite abelian group. Then $G \cong C_{n_1} \oplus \dots \oplus C_{n_r}$ where $r \in \mathbb{N}_0$, $n_1, \dots, n_r \in \mathbb{N}$ with $1 < n_1 \mid \dots \mid n_r$. We call $r = r(G)$ the *rank* of G (thus $r(G)$ is the maximum of the p -ranks of G), and a tuple (e_1, \dots, e_s) of nonzero elements of G is said to be a *basis* of G if $G = \langle e_1 \rangle \oplus \dots \oplus \langle e_s \rangle$. We start with a couple of remarks on abstract monoids, continue with the monoid of zero-sum sequences, and then we deal with Krull monoids.

By a *monoid*, we mean a commutative semigroup with identity which satisfies the cancellation law (that is, if a, b, c are elements of the monoid with $ab = ac$, then $b = c$ follows). The multiplicative semigroup of non-zero elements of an integral domain is a monoid. Let H be a monoid. We denote by H^\times the group of invertible elements of H and by $\mathcal{A}(H)$ the set of atoms (irreducible elements) of H . If $a = u_1 \cdot \dots \cdot u_k$, where $k \in \mathbb{N}$ and $u_1, \dots, u_k \in \mathcal{A}(H)$, then k is called the length of the factorization and $L(a) = \{k \in \mathbb{N} \mid a \text{ has a factorization of length } k\} \subseteq \mathbb{N}$ is the set of lengths of a . For convenience, we set $L(a) = \{0\}$ if $a \in H^\times$. Furthermore, we denote by

$$\mathcal{L}(H) = \{L(a) \mid a \in H\} \quad \text{the system of sets of lengths of } H.$$

Let $k \in \mathbb{N}$ and suppose that $H \neq H^\times$. Then

$$\mathcal{U}_k(H) = \bigcup_{a \in H, k \in \mathcal{L}(a)} \mathcal{L}(a)$$

is the union of all sets of lengths containing k . Thus, $\mathcal{U}_k(H)$ is the set of all $m \in \mathbb{N}$ such that there are atoms $u_1, \dots, u_k, v_1, \dots, v_m$ with $u_1 \cdot \dots \cdot u_k = v_1 \cdot \dots \cdot v_m$, and we define $\rho_k(H) = \sup \mathcal{U}_k(H)$. Sets of lengths are the best investigated invariants in Factorization Theory (for an overview we refer to [14, 6]).

Let G be an additively written finite abelian group. By a *sequence* over G , we mean a finite sequence of terms from G where repetition is allowed and the order is disregarded. As usual (see [14, 17]), we consider sequences as elements of the free abelian monoid $\mathcal{F}(G)$ with basis G . A sequence S over G will be written in the form

$$S = g_1 \cdot \dots \cdot g_l = \prod_{g \in G} g^{\mathbf{v}_g(S)} \in \mathcal{F}(G),$$

and we call

$$\begin{aligned} |S| &= l = \sum_{g \in G} \mathbf{v}_g(S) \in \mathbb{N}_0 \text{ the length of } S, \\ \text{supp}(S) &= \{g \in G \mid \mathbf{v}_g(S) > 0\} \subseteq G \text{ the support of } S, \text{ and} \\ \sigma(S) &= \sum_{i=1}^l g_i = \sum_{g \in G} \mathbf{v}_g(S)g \in G \text{ the sum of } S. \end{aligned}$$

We say that S is a *zero-sum sequence* if $\sigma(S) = 0$, and clearly the set of zero-sum sequences

$$\mathcal{B}(G) = \{S \in \mathcal{F}(G) \mid \sigma(S) = 0\} \subset \mathcal{F}(G)$$

is a submonoid of $\mathcal{F}(G)$, called the *monoid of zero-sum sequences* over G . Clearly, an element $A \in \mathcal{B}(G)$ is irreducible if and only if it is a minimal zero-sum sequence, and we denote by $\mathcal{A}(G) := \mathcal{A}(\mathcal{B}(G))$ the set of atoms of $\mathcal{B}(G)$. This set is finite, and the *Davenport constant* $D(G)$ of G is the maximal length of a minimal zero-sum sequence over G , thus

$$D(G) = \max\{|U| \mid U \in \mathcal{A}(G)\} \in \mathbb{N}.$$

In other words, $D(G)$ is the smallest integer ℓ such that every sequence S over G of length $|S| \geq \ell$ has a nontrivial zero-sum subsequence.

A monoid H is a *Krull monoid* if one of the following equivalent conditions is satisfied:

- (a) H is completely integrally closed and satisfies the ascending chain condition on divisorial ideals.
- (b) There is a free abelian monoid F and a homomorphism $\varphi: H \rightarrow F$ with the following property: if $a, b \in H$ and $\varphi(a)$ divides $\varphi(b)$ in F , then a divides b in H .

We refer to the monographs [18, 14] for a detailed exposition of Krull monoids and to the already mentioned paper [13]. We just mention that a domain R is a Krull domain if and only if its monoid of nonzero elements is a Krull monoid, and for monoids of modules which are Krull we refer to [2, 1, 9]. Property (a) easily shows that every integrally closed noetherian domain is a Krull domain. Since the embedding $\mathcal{B}(G) \hookrightarrow \mathcal{F}(G)$ satisfies Property (b), we infer that $\mathcal{B}(G)$ is a Krull monoid. It is easy to verify that the class group of $\mathcal{B}(G)$ is isomorphic to G and that every class contains a prime divisor. Furthermore, $\mathcal{B}(G)$ plays a universal role in the study of the arithmetic of general Krull monoids. In particular, the system of sets of lengths of a Krull monoid H with class group G , where each class contains a prime divisor, coincides with the system of sets of lengths of $\mathcal{B}(G)$. We give a precise formulation of this well-known fact (for progress in this directions see [13, Proposition 2.2]).

Proposition 2.1 ([14], Theorem 3.4.10). *Let H be a Krull monoid with class group G such that every class contains a prime divisor. Then there is a transfer homomorphism $\beta: H \rightarrow \mathcal{B}(G)$ which implies that, for every $k \in \mathbb{N}$,*

$$\mathcal{U}_k(H) = \mathcal{U}_k(\mathcal{B}(G)) \quad \text{and} \quad \rho_k(H) = \rho_k(\mathcal{B}(G)).$$

Thus the invariants $\rho_k(H)$ can be studied in the monoid of zero-sum sequences $\mathcal{B}(G)$. As usual, we set $\mathcal{U}_k(G) = \mathcal{U}_k(\mathcal{B}(G))$ and $\rho_k(G) = \rho_k(\mathcal{B}(G))$.

3. PROOF OF THEOREM 1.1

Throughout this section, let G be a finite abelian group. If $|G| \leq 2$, then $\mathcal{B}(G)$ is factorial whence $\rho_k(G) = k$ for every $k \in \mathbb{N}$. Clearly, $D^*(G) = 3$ if and only if G is cyclic of order three or isomorphic to $C_2 \oplus C_2$. In this case, Inequality (1.1) is an equality, and in particular Theorem 1.1 holds with $k^* = 1$. Thus for the remainder of this section we suppose that $D^*(G) \geq 4$, and this implies that $|G| \geq 4$.

We introduce the main concept of the present paper.

Definition 3.1. Let $A \in \mathcal{B}(G)$.

1. We say that A is *pair-nice* (with respect to G) if there is a factorization $A = U_1 \cdots U_{2k}$, $k \in \mathbb{N}$ with the following properties
 - (a) $U_1, \dots, U_{2k} \in \mathcal{A}(G)$ with $|U_1| = \dots = |U_{2k}| = D^*(G)$;
 - (b) For all $i \in [1, 2k]$, there is a $g_i \in \text{supp}(U_i)$ such that $g_1 \cdots g_{2k}$ is a product of length 2 atoms.
2. We say that A is *nice* (with respect to G) if there is a factorization $A = U_1 \cdots U_{2k+1}$, $k \in \mathbb{N}$ with the following properties
 - (a) $U_1, \dots, U_{2k+1} \in \mathcal{A}(G)$ with $|U_1| = \dots = |U_{2k+1}| = D^*(G)$;
 - (b) For all $i \in [1, 2k+1]$, there is a $g_i \in \text{supp}(U_i)$ such that one of the following holds:
 - (i) $D^*(G)$ is odd, $A(g_1 \cdots g_{2k+1})^{-1}$ is a product of length 2 atoms, and $g_1 \cdots g_{2k+1} = W_0 \cdot W_1 \cdots W_{k-1}$, where $|W_0| = 3$, $|W_1| = \dots = |W_{k-1}| = 2$, and $W_0, \dots, W_{k-1} \in \mathcal{A}(G)$.
 - (ii) $D^*(G)$ is even and there exists a $g_{2k+2} \in \text{supp}(A(g_1 \cdots g_{2k+1})^{-1})$ such that $A(g_1 \cdots g_{2k+2})^{-1}$ and $g_1 \cdots g_{2k+2}$ are both products of length 2 atoms.

Suppose there exists a *nice* $A \in \mathcal{B}(G)$, and let all notation be as in the above definition. Then

$$\{2k+1, kD^*(G) + \lfloor \frac{D^*(G)}{2} \rfloor\} \subseteq L(A) \quad \text{and hence} \quad \rho_{2k+1}(G) \geq kD^*(G) + \lfloor \frac{D^*(G)}{2} \rfloor.$$

Thus, up to a small calculation (which will be done in the actual proof of Theorem 1.1), the assertion of the theorem follows. Therefore the main task of the paper is to find nice elements. We do this for groups of rank two (Lemma 3.3), for groups of rank three (Lemma 3.4), and then we put all together in Lemma 3.5. Note, if G is cyclic of order greater than or equal to four, then there are no nice elements. Furthermore, if A is nice or pair-nice, then $0 \notin \text{supp}(A)$.

Our first lemma gathers some basic facts which we will use without further mention.

Lemma 3.2. Let E, E_1 be pair-nice zero-sum sequences (with respect to G). Suppose that $X_1, X_2, X_3 \in \mathcal{A}(G)$ are of length $D^*(G)$. Then

1. $E \cdot E_1$ is pair-nice (with respect to G);
2. If $D^*(G)$ is even and $E \cdot X_1$ is a product of length 2 atoms, then $E \cdot X_1$ is nice (with respect to G);
3. If $D^*(G)$ is odd and there exists $a_i \in \text{supp}(X_i)$ for each $i \in [1, 3]$ such that $a_1 a_2 a_3 \in \mathcal{A}(G)$ and $EX_1 X_2 X_3 (a_1 a_2 a_3)^{-1}$ is a product of length 2 atoms, then $EX_1 X_2 X_3$ is nice (with respect to G).

Proof. Since E is pair-nice, we assume that $E = U_1 \cdots U_{2k}$, where $k \in \mathbb{N}$ and $U_1, \dots, U_{2k} \in \mathcal{A}(G)$ are of length $D^*(G)$, and there exists $g_i \in \text{supp}(U_i)$ for each $i \in [1, 2k]$ such that $g_1 \cdots g_{2k}$ is a product of length 2 atoms.

1. It is obvious by definition.
2. Since $E \cdot X_1$ and $g_1 \cdots g_{2k}$ are both products of length 2 atoms, we obtain that $E(g_1 \cdots g_{2k})^{-1} X_1$ is a product of length 2 atoms. Therefore there exist $x \in \text{supp}(X_1)$ and $y \in \text{supp}(E(g_1 \cdots g_{2k})^{-1})$ such that $xy \in \mathcal{A}(G)$. It follows that $g_1 \cdots g_{2k} \cdot xy$ and $EX_1(g_1 \cdots g_{2k} \cdot xy)^{-1}$ are both products of length 2 atoms which implies that $E \cdot X_1$ is nice by $D^*(G)$ is even.

3. Since $EX_1X_2X_3(a_1a_2a_3)^{-1}$ and $g_1 \dots g_{2k}$ are product of length 2 atoms, we have that $EX_1X_2X_3(g_1 \dots g_{2k}a_1a_2a_3)^{-1}$ is a product of length 2 atoms. Moreover, $a_1a_2a_3$ is an atom implies that $EX_1X_2X_3$ is *nice* by $D^*(G)$ is odd.

□

Lemma 3.3. *Let $G = C_n \oplus C_{mn}$ with $n > 1$ and $m \in \mathbb{N}$. Then there exist a $k^* \in \mathbb{N}$ and atoms $W_1, \dots, W_{2k^*+1} \in \mathcal{A}(G)$ of length $D^*(G)$ such that $W_1 \dots W_{2k^*+1}$ is *nice*.*

Proof. Let (e_1, e_2) be a basis of G with $\text{ord}(e_1) = n$ and $\text{ord}(e_2) = mn$. Then $D^*(G) = mn + n - 1$.

Now set

$$U_i = e_1^{n-1} ((-1)^{i+1} e_2 + (i+1)e_1) ((-1)^{i+1} e_2 - ie_1) ((-1)^{i+1} e_2)^{mn-2},$$

$$V_j = e_2^{mn-1} ((-1)^{j+1} e_1 + (j+1)e_2) ((-1)^{j+1} e_1 - je_2) ((-1)^{j+1} e_1)^{n-2},$$

$$W_j = (e_1 + e_2)^{mn-1} ((-1)^{j+1} e_1 + (j+1)(e_1 + e_2)) ((-1)^{j+1} e_1 - j(e_1 + e_2)) ((-1)^{j+1} e_1)^{n-2},$$

where $i \in [0, n-1]$ and $j \in [0, mn-1]$. Then $|U_i| = |V_j| = |W_j| = D^*(G)$ and $U_i, V_j, W_j \in \mathcal{A}(G)$ for all $i \in [0, n-1]$, $j \in [0, mn-1]$.

We distinguish the following three cases.

Case 1: n is odd and m is even.

Let $n = 2\alpha + 1$ with $\alpha \geq 1$. Then $m \geq 2$ and $D^*(G)$ is even.

Since mn is even, let $E_2 = V_0 \dots V_{mn-1}$ and hence E_2 is *pair-nice*. By calculation, we obtain that $E_2 e_2^{-(mn-1)mn}$ is a product of length 2 atoms.

Let $E_3 = V_0 \dots V_{(m-1)n-1} W_{(m-1)n} \dots W_{mn-1}$ and hence E_3 is *pair-nice*. By calculation, we obtain that $E_3 \left(e_2^{(mn-1)(m-1)n} (e_1 + e_2)^{(mn-1)n} \right)^{-1}$ is a product of length 2 atoms.

Replacing the basis (e_1, e_2) with $(-e_1, e_2)$, we can construct a zero-sum sequence E'_3 similar with E_3 such that E'_3 is *pair-nice* and $E'_3 \left(e_2^{(mn-1)(m-1)n} (-e_1 + e_2)^{(mn-1)n} \right)^{-1}$ is a product of length 2 atoms.

Let $X = e_2^{mn-1} (e_1 + e_2)^{\alpha+1} (e_1 - e_2)^\alpha$ and $Y = e_2^{mn-1} (e_2 - e_1) (-e_1)^{n-1}$. Then X and Y are atoms of length $D^*(G)$. Since XY and $X(-Y)$ are *pair-nice*, we obtain that

$$E' = E_2^{2\alpha} E_3^{\alpha+1} (-E'_3)^\alpha ((-X)Y)^{\frac{(mn-1)n-1}{2}} ((-X)(-Y))^{\frac{(mn-1)n-1}{2}} \text{ is } \textit{pair-nice}.$$

By calculation we have that $(-X)E'$ is a product of length 2 atoms. It follows that $(-X)E'$ is *nice* by Lemma 3.2.2.

Case 2: n is odd and m is odd.

Then $D^*(G)$ is odd. Since n is odd, let $O_1 = U_0 \dots U_{n-1}$ and hence by calculation, we can obtain that $O_1 \left(e_1^{n(n-1)} (-e_2)^{mn} \right)^{-1}$ is a product of length 2 atoms.

Since mn is odd, let $O_2 = V_0 \dots V_{mn-1}$ and hence by calculation, $O_2 \left(e_2^{(mn-1)mn} (-e_1)^n \right)^{-1}$ is a product of length 2 atoms.

Replacing the basis (e_1, e_2) with $(-e_1, e_2)$, we can construct a zero-sum sequence O'_1 similarly with O_1 such that $O'_1 \left((-e_1)^{n(n-1)} (-e_2)^{mn} \right)^{-1}$ is a product of length 2 atoms. By the constructions of O_1 and O'_1 , we can obtain that $O_1 O'_1$ is *pair-nice* and $O_1 O'_1 (-e_2)^{-2mn}$ is a product of length 2 atoms. We denote $O_1 O'_1$ by E .

Let $X = (-e_1)^{n-1} e_2^{mn-1} (e_2 - e_1)$ and hence X is an atom of length $D^*(G)$. Therefore we let $O = O_2^{n-2} O_1 X E^{\frac{(mn-1)(n-2)}{2}}$ and hence $O \left((e_2 - e_1) e_1 (-e_2) \right)^{-1}$ is a product of length 2 atoms.

Since $O_2 U_i$ is *pair-nice* for each $i \in [0, n-1]$, we obtain that $O_2^{n-2} O_1 (U_0 U_1)^{-1}$ is *pair-nice* and hence $O(U_0 U_1 X)^{-1}$ is *pair-nice*.

By $(e_2 - e_1) \in \text{supp}(X)$, $-e_2 \in \text{supp}(U_0)$, and $e_1 \in \text{supp}(U_1)$, Lemma 3.2.3 implies that O is *nice*.

Case 3: n is even.

Then $D^*(G)$ is odd. Since n is even, let $E_1 = U_0 \cdots U_{n-1}$ and hence E_1 is *pair-nice* and $E_1 e_1^{-n(n-1)}$ is a product of length 2 atoms. Let $E_2 = V_0 \cdots V_{mn-1}$ and hence E_2 is *pair-nice* and $E_2 e_2^{-(mn-1)mn}$ is a product of length 2 atoms.

Let

$$\begin{aligned} X_1 &= e_1^{n-1} e_2^{mn-1} (e_1 + e_2), \\ X_2 &= e_2^{mn-1} e_1 (e_1 + e_2)^{\frac{n}{2}} (e_1 - e_2)^{\frac{n}{2}-1}, \\ Y &= e_1^{n-1} (-e_2)^{mn-1} (e_1 - e_2). \end{aligned}$$

Then X_1, X_2 , and Y are atoms of length $D^*(G)$. Since $X_1 Y$ and $X_1(-Y)$ are both *pair-nice*, we obtain that $(X_1 Y)^{\frac{mn}{2}} (X_1(-Y))^{\frac{mn}{2}} (-E_1)^m (-E_2)$ is *pair-nice* and

$$(X_1 Y)^{\frac{mn}{2}} (X_1(-Y))^{\frac{mn}{2}} (-E_1)^m (-E_2) (e_1 + e_2)^{-mn} \text{ is product of length 2 atoms.}$$

Replacing the basis (e_1, e_2) with $(e_1, e_2 - e_1)$, we can construct a zero-sum sequence E similarly such that E is *pair-nice* and $E e_2^{-mn}$ is a product of length 2 atoms.

Then let $E' = (X_1 Y)^{\frac{n}{2}} (X_1(-Y))^{\frac{n}{2}} (-E_1) (-E)^n$ and hence E' is *pair-nice* and

$$E' ((e_1 + e_2)^n (-e_2)^n)^{-1} \text{ is a product of length 2 atoms.}$$

Similarly with E' , if we replace the basis (e_1, e_2) with $(e_1, -e_2)$, we can construct a zero-sum sequence E'' such that E'' is *pair-nice* and

$$E'' ((e_1 - e_2)^n e_2^n)^{-1} \text{ is a product of length 2 atoms.}$$

Since $X_2 Y$ and $X_2(-Y)$ are both *pair-nice*, we let $E''' = (X_2 Y)^{\frac{n}{2}} (X_2(-Y))^{\frac{n}{2}} (-E)^n (-E')^{\frac{n}{2}} (-E'')^{\frac{n}{2}-1}$ and hence E''' is *pair-nice* and $E''' e_1^{-n}$ is a product of length 2 atoms. It follows that $(-E)(-E''')$ is *pair-nice*, $(e_1 + e_2) \in \text{supp}(X_1)$, $-e_2 \in \text{supp}(Y)$, $-e_1 \in \text{supp}(-Y)$, and

$$X_1 Y (-Y) (-E) (-E''') ((-e_2)(-e_1)(e_1 + e_2))^{-1} \text{ is a product of length 2 atoms}$$

which implies that $X_1 Y (-Y) (-E) (-E''')$ is *nice* by Lemma 3.2.3. \square

Lemma 3.4. *Let $G = C_{n_1} \oplus C_{n_2} \oplus C_{n_3}$ with $1 < n_1 | n_2 | n_3$. Then there exist a $k^* \in \mathbb{N}$ and atoms $W_1, \dots, W_{2k^*+1} \in \mathcal{A}(G)$ of length $D^*(G)$ such that $W_1 \cdots W_{2k^*+1}$ is nice.*

Proof. Let (e_1, e_2, e_3) be a basis of G with $\text{ord}(e_1) = n_1$, $\text{ord}(e_2) = n_2$, and $\text{ord}(e_3) = n_3$. Then $D^*(G) = n_1 + n_2 + n_3 - 2$. Denote

$$\begin{aligned} X_1 &= e_1^{n_1-1} e_2^{n_2-1} (-e_3)^{n_3-2} (e_1 - e_3)(e_2 - e_3), \\ X_2 &= e_1^{n_1-1} (-e_2)^{n_2-2} e_3^{n_3-1} (e_1 - e_2)(e_3 - e_2), \\ X_3 &= (-e_1)^{n_1-2} e_2^{n_2-1} e_3^{n_3-1} (-e_1 + e_2)(-e_1 + e_3). \end{aligned}$$

It is easy to see that X_i is an atom of length $D^*(G)$ for each $i \in [1, 3]$. Thus

$$X_1 X_2 X_3 (e_1^{n_1} e_2^{n_2} e_3^{n_3})^{-1} \text{ is a product of length 2 atoms.}$$

If $n_1 = n_2 = n_3 = 2$, we have $X_1 X_2 X_3$ is a product of length 2 atoms and hence $X_1 X_2 X_3$ is *nice*. Thus we can assume $n_3 \geq 4$.

Denote

$$\begin{aligned} X'_1 &= e_1^{n_1-1} (-e_2)^{n_2-1} e_3^{n_3-2} (e_1 + e_3)(-e_2 + e_3), \\ X'_2 &= e_1^{n_1-1} e_2^{n_2-2} (-e_3)^{n_3-1} (e_1 + e_2)(-e_3 + e_2), \\ X'_3 &= (-e_1)^{n_1-2} (-e_2)^{n_2-1} (-e_3)^{n_3-1} (-e_1 - e_2)(-e_1 - e_3). \end{aligned}$$

Thus $E_1 = X_1 X_1' X_2 X_2' X_3 X_3'$ is *pair-nice* and $E_1 e_1^{-2n_1}$ is a product of length 2 atoms. Similarly, we can construct *pair-nice* zero-sum sequences E_2 and E_3 such that $E_2 e_2^{-2n_2}$ and $E_3 e_3^{-2n_3}$ are both products of length 2 atoms.

Set

$$\begin{aligned} U_i &= e_1^{n_1-1} ((-1)^{i+1} e_3 + (i+1) e_1) ((-1)^{i+1} e_3 - i e_1) ((-1)^{i+1} e_3)^{n_3-3} ((-1)^{i+1} (e_2 + e_3)) ((-1)^{i+1} e_2)^{n_2-1}, \\ V_j &= e_2^{n_2-1} ((-1)^{j+1} e_3 + (j+1) e_2) ((-1)^{j+1} e_3 - j e_2) ((-1)^{j+1} e_3)^{n_3-3} ((-1)^{j+1} (e_1 + e_3)) ((-1)^{j+1} e_1)^{n_1-1}, \\ W_l &= \begin{cases} e_3^{n_3-1} ((-1)^{l+1} e_2 + (l+1) e_3) ((-1)^{l+1} e_2 - l e_3) ((-1)^{l+1} e_2)^{n_2-3} ((-1)^{l+1} (e_1 + e_2)) ((-1)^{l+1} e_1)^{n_1-1}, & \text{if } n_2 \geq 3, \\ e_3^{n_3-1} (e_1 + l e_2 + (l+1) e_3) (e_1 + (l+1) e_2 - l e_3) e_2, & \text{if } n_1 = n_2 = 2, \end{cases} \end{aligned}$$

where $i \in [0, n_1 - 1]$, $j \in [0, n_2 - 1]$, and $l \in [0, n_3 - 1]$. It is easy to see that, V_i , U_j , W_l are all atoms of length $D^*(G)$, where $i \in [0, n_1 - 1]$, $j \in [0, n_2 - 1]$, and $l \in [0, n_3 - 1]$.

Now we distinguish the following four cases.

Case 1: n_1 is even.

Then $D^*(G)$ is even since $1 < n_1 \mid n_2 \mid n_3$.

Since n_1 is even, we let $E_1' = U_0 \cdots U_{n_1-1}$, $E_2' = V_0 \cdots V_{n_2-1}$, and $E_3' = W_0 \cdots W_{n_3-1}$ and hence E_1' , E_2' , and E_3' are *pair-nice*. Moreover $E_1' e_1^{-n_1(n_1-1)}$, $E_2' e_2^{-n_2(n_2-1)}$, and $E_3' e_3^{-n_3(n_3-1)}$ are products of length 2 atoms.

Therefore let $E_4 = E_1' (-E_1)^{\frac{n_1}{2}-1}$, $E_5 = E_2' (-E_2)^{\frac{n_2}{2}-1}$, and $E_6 = E_3' (-E_3)^{\frac{n_3}{2}-1}$, and hence E_4, E_5, E_6 are *pair-nice* and $E_4 e_1^{-n_1}$, $E_5 e_2^{-n_2}$, and $E_6 e_3^{-n_3}$ are products of length 2 atoms.

It follows that $O = E_4 E_5 E_6 (-X_1) (-X_2) (-X_3)$ is a product of length 2 atoms and $E_4 E_5 E_6 (-X_1) (-X_2)$ is *pair-nice*. Then O is *nice* by Lemma 3.2.2.

Case 2: n_1 is odd, n_2 is even.

Then $D^*(G)$ is odd. Since n_1 is odd, we let $O_1 = U_0 \cdots U_{n_1-1}$ and hence $O_1 U_0^{-1}$ is *pair-nice* and

$$O_1 \left(e_1^{(n_1-1)n_1} (-e_3)^{n_3-1} (-e_2 - e_3) (-e_2)^{n_2-1} \right)^{-1} \text{ is a product of length 2 atoms.}$$

Since n_2 is even, we let $E_2' = V_0 \cdots V_{n_2-1}$ and $E_3' = W_0 \cdots W_{n_3-1}$ and hence E_2' and E_3' are *pair-nice*. Moreover $E_2' e_2^{-n_2(n_2-1)}$ and $E_3' e_3^{-n_3(n_3-1)}$ are both products of length 2 atoms.

Therefore let $E_5 = E_2' (-E_2)^{\frac{n_2}{2}-1}$ and $E_6 = E_3' (-E_3)^{\frac{n_3}{2}-1}$, and hence E_5, E_6 are *pair-nice* and $E_5 e_2^{-n_2}$ and $E_6 e_3^{-n_3}$ are both products of length 2 atoms.

Let $O = (-E_1)^{\frac{n_1-1}{2}} E_5 E_6 O_1 X_1 (-X_1)$ and hence $O (e_2 e_3 (-e_2 - e_3))^{-1}$ is a product of length 2 atoms.

Since $(-e_2 - e_3) \in \text{supp}(U_0)$, $e_2 \in \text{supp}(X_1)$, $e_3 \in \text{supp}(-X_1)$, and $O (U_0 X_1 (-X_1))^{-1}$ is *pair-nice*, we obtain that O is *nice* by Lemma 3.2.3.

Case 3: n_1 is odd, n_2 is odd, and n_3 is even.

Then $D^*(G)$ is even. Since n_1 is odd, we let $O_1 = U_0 \cdots U_{n_1-1}$ and hence $O_1 U_0^{-1}$ is *pair-nice* and

$$O_1 \left(e_1^{(n_1-1)n_1} (-e_3)^{n_3-1} (-e_2 - e_3) (-e_2)^{n_2-1} \right)^{-1} \text{ is a product of length 2 atoms.}$$

Since n_3 is even, we let $E_3' = W_0 \cdots W_{n_3-1}$ and hence E_3' is *pair-nice* and $E_3' e_3^{-(n_3-1)n_3}$ is a product of length 2 atoms. Therefore $E_6 = E_3' (-E_3)^{\frac{n_3}{2}-1}$ is *pair-nice* and $E_6 e_3^{-n_3}$ is a product of length 2 atoms.

Since $n_2 \geq 3$, we let

$$\begin{aligned} W_l' &= (e_2 + e_3)^{n_3-1} ((-1)^{l+1} e_2 + (l+1)(e_2 + e_3)) ((-1)^{l+1} e_2 - l(e_2 + e_3)) ((-1)^{l+1} e_2)^{n_2-3} \\ &\quad ((-1)^{l+1} (e_1 + e_2)) ((-1)^{l+1} e_1)^{n_1-1}, \end{aligned}$$

where $l \in [0, n_3 - 1]$. Thus W_l' is an atom of length $D^*(G)$ for each $l \in [0, n_3 - 1]$.

Let $E_7 = (W_0 W_1) \cdots (W_{n_3-n_2-1} W'_{n_3-n_2}) \cdots (W'_{n_3-2} W'_{n_3-1})$ and hence E_7 is *pair-nice* and

$$E_7 \left(e_3^{(n_3-1)(n_3-n_2)} (e_2 + e_3)^{(n_3-1)n_2} \right)^{-1} \text{ is a product of length 2 atoms .}$$

Therefore let $O' = O_1^{(n_3-1)n_2} E_7 (-E_1)^{\frac{(n_3-1)n_2(n_1-1)n_1}{2n_1}} E_2^{\frac{(n_3-1)n_2(n_2-1)}{2n_2}} E_6^{\frac{(n_3-1)n_3(n_2-1)}{n_3}}$ and hence O' is a product of length 2 atoms and $O'(U_0)^{-(n_3-1)n_2}$ is *pair-nice*.

Let $Y = e_1^{n_1-1} e_2^{n_2-1} e_3^{n_3-1} (e_1 + e_2 + e_3)$ and hence Y is an atom of length $D^*(G)$. Since YU_0 and $(-Y)U_0$ are both *pair-nice*, we obtain that $U_0^{(n_3-1)n_2-1} (Y(-Y))^{\frac{(n_3-1)n_2-1}{2}} = (U_0 Y)^{\frac{(n_3-1)n_2-1}{2}} (U_0(-Y))^{\frac{(n_3-1)n_2-1}{2}}$ is *pair-nice*.

It follows that $O = O'(Y(-Y))^{\frac{(n_3-1)n_2-1}{2}}$ is a product of length 2 atoms and

$$OU_0^{-1} = O'(U_0)^{-(n_3-1)n_2} \cdot U_0^{(n_3-1)n_2-1} (Y(-Y))^{\frac{(n_3-1)n_2-1}{2}} \text{ is } \textit{pair-nice}.$$

Then O is *nice* by Lemma 3.2.2.

Case 4: n_1 is odd, n_2 is odd, and n_3 is odd.

Then $D^*(G)$ is odd. Since n_3 is odd, we let $O_3 = W_0 \cdots W_{n_3-1}$ and hence $O_3 W_0^{-1}$ is *pair-nice* and

$$O_3 \left(e_3^{(n_3-1)n_3} (-e_2)^{n_2-1} (-e_1 - e_2) (-e_1)^{n_1-1} \right)^{-1} \text{ is a product of length 2 atoms ,}$$

and hence

$$O_3 (-E_3)^{\frac{n_3-1}{2}} X_1 X_2 X_3 (e_3^{n_3} e_1 e_2 (-e_1 - e_2))^{-1} \text{ is a product of length 2 atoms .}$$

By $(W_0 X_1)(X_2 X_3)$ is *pair-nice*, we have that $O_3 (-E_3)^{\frac{n_3-1}{2}} X_1 X_2 X_3$ is *pair-nice*.

Similarly, if we replace the basis (e_1, e_2, e_3) with $(-e_1, -e_2, e_1 + e_2 + e_3)$, we can construct a zero-sum sequence E such that E is *pair-nice* and

$$E ((e_1 + e_2 + e_3)^{n_3} (-e_1) (-e_2) (e_1 + e_2))^{-1} \text{ is a product of length 2 atoms .}$$

Let

$$\begin{aligned} Y &= e_1^{n_1-1} e_2^{n_2-1} e_3^{n_3-1} (e_1 + e_2 + e_3), \\ Y_1 &= e_1^{n_1-1} (-e_1 - e_2)^{n_2-1} (-e_3)^{n_3-1} (-e_2 - e_3), \end{aligned}$$

and hence Y and Y_1 are atoms of length $D^*(G)$.

Therefore let

$$\begin{aligned} O_4 &= Y^{n_3} (-E_1)^{\frac{(n_1-1)n_3}{2n_1}} (-E_2)^{\frac{(n_2-1)n_3}{2n_2}} (-E_3)^{\frac{(n_3-1)}{2}}, \\ O &= O_4 (Y_1 (-Y_1))^{\frac{n_3-1}{2}} (-E). \end{aligned}$$

Then $O_4 (e_1 + e_2 + e_3)^{-n_3}$ is a product of length 2 atoms and hence $O (e_1 e_2 (-e_1 - e_2))^{-1}$ is a product of length 2 atoms. By calculation, we obtain that

$$O(Y^2 Y_1)^{-1} = O_4 (Y)^{-n_3} \cdot (Y Y_1)^{\frac{n_3-3}{2}} \cdot (Y (-Y_1))^{\frac{n_3-1}{2}} \cdot (-E) \text{ is } \textit{pair-nice},$$

Therefore $e_1 \in \text{supp}(Y)$, $e_2 \in \text{supp}(Y)$, and $-e_1 - e_2 \in \text{supp}(Y_1)$ imply that O is *nice* by Lemma 3.2.3. \square

Lemma 3.5. *Let $G = G_1 \oplus G_2$, where $G_1, G_2 \subset G$ are noncyclic subgroups of G satisfying $r(G) = r(G_1) + r(G_2)$. Suppose that there exist $k \in \mathbb{N}$ and atoms $U_1, \dots, U_{2k+1} \in \mathcal{A}(G_1)$ of length $D^*(G_1)$ and atoms $V_1, \dots, V_{2k+1} \in \mathcal{A}(G_2)$ of length $D^*(G_2)$ such that $U_1 \cdots U_{2k+1}$ is nice (with respect to G_1) and $V_1 \cdots V_{2k+1}$ is nice (with respect to G_2). Then there exist atoms $W_1, \dots, W_{2k+1} \in \mathcal{A}(G)$ of length $D^*(G)$ such that $W_1 \cdots W_{2k+1}$ is nice (with respect to G).*

Proof. Without loss of generality, we can distinguish the following three cases.

Case 1. $D^*(G_1)$ and $D^*(G_2)$ are odd.

Since $U_1 \cdots U_{2k+1}$ is *nice* (with respect to G_1), without loss of generality, we can assume that there exist $g_i \in \text{supp}(U_i)$ for each $i \in [1, 2k+1]$ such that $\sigma(g_1 g_2 g_3) = 0$, $g_{2j} = -g_{2j+1}$ for each $j \in [2, k]$, and $U_1 \cdots U_{2k+1}(g_1 \cdots g_{2k+1})^{-1}$ is a product of length 2 atoms.

With the same reason, we can assume that there exist $h_i \in \text{supp}(V_i)$ for each $i \in [1, 2k+1]$ such that $\sigma(h_1 h_2 h_3) = 0$, $h_{2j} = -h_{2j+1}$ for each $j \in [2, k]$, and $V_1 \cdots V_{2k+1}(h_1 \cdots h_{2k+1})^{-1}$ is a product of length 2 atoms.

Let $W_i = U_i g_i^{-1} \cdot V_i h_i^{-1} \cdot (g_i + h_i)$ for all $i \in [1, 2k+1]$. Then W_i is an atom over G of length $D^*(G_1) + D^*(G_2) - 1 = D^*(G)$, and

$$(g_1 + h_1) \cdots (g_{2k+1} + h_{2k+1}) = Z_0 \cdot Z_1 \cdots Z_{k-1}$$

where $Z_0 = (g_1 + h_1)(g_2 + h_2)(g_3 + h_3) \in \mathcal{A}(G)$ and $Z_i = (g_{2i+2} + h_{2i+2})(g_{2i+3} + h_{2i+3}) \in \mathcal{A}(G)$ for each $i \in [1, k-1]$,

$W_1 \cdots W_{2k+1} ((g_1 + h_1) \cdots (g_{2k+1} + h_{2k+1}))^{-1} = U_1 \cdots U_{2k+1} (g_1 \cdots g_{2k+1})^{-1} \cdot V_1 \cdots V_{2k+1} (h_1 \cdots h_{2k+1})^{-1}$ is a product of atoms of length 2.

It follows that $W_1 \cdots W_{2k+1}$ is *nice* (with respect to G) by $D^*(G)$ is odd.

Case 2. $D^*(G_1)$ and $D^*(G_2)$ are even.

Since $U_1 \cdots U_{2k+1}$ is *nice* (with respect to G_1), without loss of generality, we can assume that there exist $g_i \in \text{supp}(U_i)$ for each $i \in [1, 2k+1]$ and $g_{2k+2} \in \text{supp}(U_1 g_1^{-1})$ such that $g_1 = -g_2$, $g_{2k+2} = -g_3$, and $g_{2j} = -g_{2j+1}$ for each $j \in [2, k]$ and $U_1 \cdots U_{2k+1}(g_1 \cdots g_{2k+2})^{-1}$ is a product of length 2 atoms.

With the same reason, we can assume that there exist $h_i \in \text{supp}(V_i)$ for each $i \in [1, 2k+1]$ and $h_{2k+2} \in \text{supp}(V_1 h_1^{-1})$ such that $h_1 = -h_2$, $h_{2k+2} = -h_3$, and $h_{2j} = -h_{2j+1}$ for each $j \in [2, k]$ and $V_1 \cdots V_{2k+1}(h_1 \cdots h_{2k+2})^{-1}$ is a product of atoms of length 2.

Let $W_i = U_i g_i^{-1} \cdot V_i h_i^{-1} \cdot (g_i + h_i)$ for all $i \in [1, 2k+1]$ and

$$W_1 = U_1 (g_1 g_{2k+2})^{-1} \cdot V_2 h_2^{-1} \cdot (g_1 + h_2) \cdot g_{2k+2},$$

$$W_2 = U_2 g_2^{-1} \cdot V_1 (h_1 h_{2k+2})^{-1} \cdot (g_2 + h_1) \cdot h_{2k+2},$$

$$W_3 = U_3 g_3^{-1} \cdot V_3 h_3^{-1} \cdot (g_3 + h_3).$$

Then W_i is an atom over G of length $D^*(G_1) + D^*(G_2) - 1 = D^*(G)$ for all $i \in [1, 2k+1]$. It follows that

$$g_{2k+2} \cdot h_{2k+2} \cdot (g_3 + h_3) \cdot (g_4 + h_4) \cdots (g_{2k+1} + h_{2k+1}) = Z_0 \cdot Z_1 \cdots Z_{k-1},$$

where $Z_0 = g_{2k+2} h_{2k+2} (g_3 + h_3) \in \mathcal{A}(G)$ and $Z_i = (g_{2i+2} + h_{2i+2})(g_{2i+3} + h_{2i+3}) \in \mathcal{A}(G)$ for each $i \in [1, k-1]$.

Moreover

$$\begin{aligned} & W_1 \cdots W_{2k+1} (g_{2k+2} \cdot h_{2k+2} \cdot (g_3 + h_3) \cdot (g_4 + h_4) \cdots (g_{2k+1} + h_{2k+1}))^{-1} \\ &= U_1 \cdots U_{2k+1} (g_1 \cdots g_{2k+2})^{-1} \cdot V_1 \cdots V_{2k+1} (h_1 \cdots h_{2k+2})^{-1} \cdot (g_1 + h_2)(g_2 + h_1) \end{aligned}$$

is a product of atoms of length 2.

Thus $W_1 \cdots W_{2k+1}$ is *nice* (with respect to G) by $D^*(G)$ is odd.

Case 3. $D^*(G_1)$ is even and $D^*(G_2)$ is odd.

Since $U_1 \cdots U_{2k+1}$ is *nice* (with respect to G_1) and $D^*(G_1)$ is even, without loss of generality, we can assume that there exist $g_i \in \text{supp}(U_i)$ for each $i \in [1, 2k+1]$ and $g_{2k+2} \in \text{supp}(U_1 g_1^{-1})$ such that $g_1 = -g_2$, $g_{2k+2} = -g_3$, and $g_{2j} = -g_{2j+1}$ for each $j \in [2, k]$ and $U_1 \cdots U_{2k+1}(g_1 \cdots g_{2k+2})^{-1}$ is a product of length 2 atoms.

Since $V_1 \cdots V_{2k+1}$ is *nice* (with respect to G_2) and $D^*(G_2)$ is odd, without loss of generality, we can assume that there exist $h_i \in \text{supp}(V_i)$ for each $i \in [1, 2k+1]$ such that $\sigma(h_1 h_2 h_3) = 0$ and $h_{2j} = -h_{2j+1}$ for each $j \in [2, k]$ and $V_1 \cdots V_{2k+1}(h_1 \cdots h_{2k+1})^{-1}$ is a product of length 2 atoms.

Let $W_i = U_i g_i^{-1} \cdot V_i h_i^{-1} \cdot (g_i + h_i)$ for all $i \in [4, 2k+1]$ and

$$\begin{aligned} W_1 &= U_1(g_1 g_{2k+2})^{-1} \cdot V_1 h_1^{-1} \cdot (g_1 - h_2) \cdot (g_{2k+2} + h_1 + h_2), \\ W_2 &= U_2 g_2^{-1} \cdot V_2 h_2^{-1} \cdot (g_2 + h_2), \\ W_3 &= U_3 g_3^{-1} \cdot V_3 h_3^{-1} \cdot (g_3 + h_3). \end{aligned}$$

Then W_i is an atom over G of length $D^*(G_1) + D^*(G_2) - 1 = D^*(G)$ for all $i \in [1, 2k+1]$. It follows that

$$(g_1 - h_2) \cdot (g_{2k+2} + h_1 + h_2) \cdot (g_2 + h_2) \cdot (g_3 + h_3) \cdot (g_4 + h_4) \cdots (g_{2k+1} + h_{2k+1})$$

is a product of length 2 atoms and

$$\begin{aligned} &W_1 \cdots W_{2k+1} \left((g_1 - h_2) \cdot (g_{2k+2} + h_1 + h_2) \cdot (g_2 + h_2) \cdot (g_3 + h_3) \cdot (g_4 + h_4) \cdots (g_{2k+1} + h_{2k+1}) \right)^{-1} \\ &= U_1 \cdots U_{2k+1} (g_1 \cdots g_{2k+2})^{-1} \cdot V_1 \cdots V_{2k+1} (h_1 \cdots h_{2k+1})^{-1} \end{aligned}$$

is a product of length 2 atoms.

Thus $W_1 \cdots W_{2k+1}$ is *nice* (with respect to G) by $D^*(G)$ is even. □

Proof of Theorem 1.1. Let H be a Krull monoid with finite noncyclic class group G such that every class contains a prime divisor. By Proposition 2.1 we have $\rho_k(H) = \rho_k(G)$ for every $k \in \mathbb{N}$. If $G \cong C_2 \oplus C_2$, then $D^*(G) = 3$ and the assertion of the theorem follows from Inequality 1.1 with $k^* = 1$. Suppose that $D^*(G) \geq 4$. We start with the following assertion.

Assertion. There exist a $k^* \in \mathbb{N}$ and atoms W_1, \dots, W_{2k^*+1} over G of length $D^*(G)$ such that $W_1 \cdots W_{2k^*+1}$ is *nice* (with respect to G).

Proof of Assertion. We proceed by induction on $r(G)$.

If $r(G) = 2$ or 3 , then the Assertion follows by Lemma 3.3 and 3.4. Assume that $r(G) \geq 4$ and suppose that the Assertion is true for all groups of smaller rank. Let $G = G_1 \oplus G_2$ with $r(G_1) = r - 2$ and $r(G_2) = 2$. Then by our assumption, there exist a $k_1 \in \mathbb{N}$ and atoms U_1, \dots, U_{2k_1+1} over G_1 of length $D^*(G_1)$ such that $U_1 \cdots U_{2k_1+1}$ is *nice* (with respect to G_1). By Lemma 3.3, there exist a $k_2 \in \mathbb{N}$ and atoms V_1, \dots, V_{2k_2+1} over G_2 of length $D^*(G_2)$ such that $V_1 \cdots V_{2k_2+1}$ is *nice* (with respect to G_2).

Let $k^* = \max(k_1, k_2)$. Without loss of generality, we can assume that $k_1 = k^* \geq k_2$. Thus $k_1 - k_2$ is even and hence $V_1 \cdots V_{2k_2+1} \cdot (V_1(-V_1))^{k_1-k_2}$ is *nice* (with respect to G_2). Therefore the Assertion follows by Lemma 3.5. □(Proof of Assertion)

By the very definition of nice elements (and outlined in detail after Definition 3.1), it follows that

$$(3.1) \quad \rho_{2k^*+1}(G) \geq k^* D^*(G) + \left\lfloor \frac{D^*(G)}{2} \right\rfloor.$$

Let $k \geq k^*$. Since $\rho_{2(k-k^*)}(G) = (k - k^*)D(G)$ and $\mathcal{U}_{2(k-k^*)}(G) + \mathcal{U}_{2k^*+1}(G) \subseteq \mathcal{U}_{2k+1}(G)$, it follows that

$$\rho_{2k+1}(G) \geq \rho_{2(k-k^*)}(G) + \rho_{2k^*+1}(G) \geq (k - k^*)D(G) + k^* D^*(G) + \left\lfloor \frac{D^*(G)}{2} \right\rfloor.$$

If $D(G) = D^*(G)$, then the assertion follows from Inequality 1.1. □

REFERENCES

1. N.R. Baeth and A. Geroldinger, *Monoids of modules and arithmetic of direct-sum decompositions*, Pacific J. Math. **271** (2014), 257 – 319.
2. N.R. Baeth and R. Wiegand, *Factorization theory and decomposition of modules*, Am. Math. Mon. **120** (2013), 3 – 34.
3. P. Baginski, S.T. Chapman, N. Hine, and J. Paixao, *On the asymptotic behavior of unions of sets of lengths in atomic monoids*, Involve, a journal of mathematics **1** (2008), 101 – 110.
4. V. Blanco, P. A. García-Sánchez, and A. Geroldinger, *Semigroup-theoretical characterizations of arithmetical invariants with applications to numerical monoids and Krull monoids*, Illinois J. Math. **55** (2011), 1385 – 1414.
5. G. Bhowmik and J.-C. Schlage-Puchta, *Davenport’s constant for groups of the form $\mathbb{Z}_3 \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_{3d}$* , in *Additive Combinatorics*, CRM Proceedings and Lecture Notes **43**, 307 – 326, American Mathematical Society (2007).
6. S.T. Chapman and M. Fontana and A. Geroldinger and B. Olberding, *Multiplicative Ideal Theory and Factorization Theory*, Springer, to appear.
7. S.T. Chapman and W.W. Smith, *Factorization in Dedekind domains with finite class group*, Isr. J. Math. **71** (1990), 65 – 95.
8. A. Czogala, *Arithmetic characterization of algebraic number fields with small class number*, Math. Z. **176** (1981), 247 – 253.
9. A. Facchini, *Krull monoids and their application in module theory*, Algebras, Rings and their Representations (A. Facchini, K. Fuller, C. M. Ringel, and C. Santa-Clara, eds.), World Scientific, 2006, pp. 53 – 71.
10. M. Freeze and A. Geroldinger, *Unions of sets of lengths*, Funct. Approximatio, Comment. Math. **39** (2008), 149 – 162.
11. W. Gao and A. Geroldinger, *On products of k atoms*, Monatsh. Math. **156** (2009), 141 – 157.
12. A. Geroldinger, *Additive group theory and non-unique factorizations*, Combinatorial Number Theory and Additive Group Theory (A. Geroldinger and I. Ruzsa, eds.), Advanced Courses in Mathematics CRM Barcelona, Birkhäuser, 2009, pp. 1 – 86.
13. A. Geroldinger, D.J. Gryniewicz, and P. Yuan, *On products of k atoms II*, Mosc. J. Comb. Number Theory, to appear, arXiv:1503.06164.
14. A. Geroldinger and F. Halter-Koch, *Non-Unique Factorizations. Algebraic, Combinatorial and Analytic Theory*, Pure and Applied Mathematics, vol. 278, Chapman & Hall/CRC, 2006.
15. A. Geroldinger, F. Kainrath, and A. Reinhart, *Arithmetic of seminormal weakly Krull monoids and domains*, J. Algebra, to appear.
16. A. Geroldinger, M. Liebmann, and A. Philipp, *On the Davenport constant and on the structure of extremal sequences*, Period. Math. Hung. **64** (2012), 213 – 225.
17. D.J. Gryniewicz, *Structural Additive Theory*, Springer (2013).
18. F. Halter-Koch, *Ideal Systems. An Introduction to Multiplicative Ideal Theory*, Marcel Dekker, 1998.
19. L. Salce and P. Zanardo, *Arithmetical characterization of rings of algebraic integers with cyclic ideal class group*, Boll. Unione. Mat. Ital., VI. Ser., D, Algebra Geom. **1** (1982), 117 – 122.

MATHEMATICAL COLLEGE, CHINA UNIVERSITY OF GEOSCIENCES (BEIJING), HAIDIAN DISTRICT, BEIJING, THE PEOPLE’S REPUBLIC OF CHINA

E-mail address: fys@cugb.edu.cn

INSTITUTE FOR MATHEMATICS AND SCIENTIFIC COMPUTING, UNIVERSITY OF GRAZ, NAWI GRAZ, HEINRICHSTRASSE 36, 8010 GRAZ, AUSTRIA

E-mail address: qinghai.zhong@uni-graz.at